

Assessment of Thyroid Function in Pregnant Females Attending Suez Canal University Hospital

Original
Article

A. A. Abo El-Roose¹, N. H. Al-Imam², A. M. El-Deib², G. A. Tawfik²

¹Department of Obstetrics and Gynecology, ²Department of Internal Medicine, Faculty of Medicine, Suez-Canal University, Egypt

ABSTRACT

Background: Pregnancy is a physiological state in which significant changes in thyroid function occur. Several factors contribute to these changes. These factors could contribute to thyroid dysfunction during pregnancy especially when a deficiency of iodine intake exists and when thyroid reserve is not sufficient.

Aim: To study thyroid functions in pregnant women avoiding maternal and fetal complications associated with thyroid dysfunctions.

Materials and Methods: A cross-sectional study was carried out on 100 pregnant women attending Obstetrics Outpatient Clinic in Suez-Canal University Hospitals were invited to enroll in the study. At the end of study, the blood samples were assessed for free T3, free T4 and TSH.

Results: This study revealed that most of the pregnant women had normal thyroid functions (51%), while subclinical hypothyroidism (39%) was the most prevalent disorder followed by clinical hypothyroidism (6%) and isolated hypothyroxinemia (4%).

Conclusion: The most prevalent pattern of thyroid dysfunction in pregnant women was subclinical hypothyroidism.

Key Words: Thyroid, TSH, T3, T4, pregnancy

Received: 29 September 2019, **Accepted:** 29 September 2019

Corresponding Author: Ahmed Abdel-Monem Abo El-Roose, Department of Obstetrics and Gynecology, Faculty of Medicine, Suez-Canal University, Egypt, **Tel.:** 01223562341, **E-mail:** aaboelroose@gmail.com

ISSN: 2090-7265, August 2020, Vol.10, No. 3

INTRODUCTION

Thyroid dysfunction is the second most common endocrine disorder, only after diabetes mellitus, affecting females in reproductive age group. Thyroid gland performs its very important functions. These hormones also play a very important role in maturation of fetal brain cells^[1].

Pregnancy is a physiological state in which significant changes in thyroid function occur. Several factors contribute to these changes. Indeed, since the beginning of pregnancy, the HCG (human chorionic gonadotrophin) secreted by the placenta, given its homology of structure with TSH (thyroid stimulating hormone), exerts a stimulatory effect on thyroid gland leading to an increase in the secretion of thyroid hormones (T3 and T4) and a decrease in TSH, especially during the first trimester^[2].

In addition, pregnancy induces increase in metabolic function and requires a higher production of thyroid hormone. Demand for iodine intake increases especially as there is a greater urinary excretion of iodine and a transfer of iodine to the fetus when his thyroid becomes functional^[3].

All these factors could contribute to thyroid dysfunction during pregnancy especially when a deficiency of iodine intake exists and when thyroid reserve is not sufficient.

PATIENTS AND METHODS

Study design: The present study was descriptive cross-section study. The study was carried out in Suez-Canal University Hospitals, Obstetrics Outpatient Clinic, Ismailia, Egypt.

Study population: A total number of one hundred pregnant women attending Obstetrics Outpatient Clinic at Ismailia University Hospitals for antenatal care were invited to enroll in this study. The included patients had the following criteria: age 18-40 years old pregnant females, single embryo, first, second and third trimester of pregnancy. Patients were excluded if they were smokers, patients on medications that affect thyroid profile as amiodarone, patients refuse to be included in the study, women with a family history of thyroid disease, women with a history of either miscarriage or preterm delivery, women with certain other autoimmune disorders which are frequently associated with autoimmune thyroid dysfunction

as vitiligo, adrenal insufficiency etc. and women with prior therapeutic head or neck irradiation.

Tools and data collection : Patients were assessed through personal interview. The assessment included history taking, general and neck examination. All patients had thyroid function test included TSH, FT3, FT4. As no trimester-specific reference ranges for the assessment of thyroid function in pregnancy had been established for the Egyptian population at the time of the study, the Guidelines of the American Thyroid Association (ATA) for the Diagnosis and Management of Thyroid Disease during Pregnancy and Postpartum recommendations were applied^[4].

STATISTICAL ANALYSIS:

Collected data was processed using SPSS 18. Quantitative data were presented as mean \pm SD while qualitative data were expressed as numbers and percentages. Unpaired T-test was used to test the significance of difference between quantitative data while Chi-Square test was used for qualitative data. A probability value (*P-value*) less than 0.05 was considered significant. The results were presented in tables and graphs.=

RESULTS

In the present study, various thyroid disorders among one hundred (N=100) pregnant women attending Obstetric Outpatient Clinic in Suez-Canal University Hospital were assessed.

The demographic data of the studied population was demonstrated. It showed that the study population age

ranged from 18 to 39 years with average of 28.5 ± 5 years. Sixty percent of the studied population was younger than thirty years old. Seventy two women were residents in urban areas and 81% of the studied population was multigravida. Most of the studied population was in the first trimester representing 45% of all (Table 1).

Chart 1 showed that 49% of the studied population had thyroid disorder. Subclinical hypothyroidism was the most prevalent disorder representing 39% of the total population.

The highest prevalence of hypothyroidism was in the second trimester (50%) followed by first trimester (48.9%) and finally third trimester (47.6%) which showed that no statistically significant relationship between each trimester of pregnancy and development of thyroid disorders (Table 2).


Concerning the relation between thyroid function and age, 3 (50%) of patients with clinical hypothyroidism were less than 30 years and 3 (50%) were 30-40 years old and no statistically significant relationship between thyroid functions and age was revealed (Table 3).

According to ATA guide-lines 2017, the thyroid function laboratory results of the enrolled pregnant women were 77% with a normal thyroid function, 13% diagnosed to have subclinical hypothyroidism, 3% clinical hypothyroidism and 7% isolated hypothyroxinemia (Table 4).

So, there is over estimation of thyroid dysfunction using ATA guide lines 2017 and under estimation using only selected high risk population screening.

Table 1 : The demographic data of the studied population

Demographic characteristics	No.	%
Age		
18-29	60	60.0
≥30	40	40.0
Min. – Max.		18.0 – 39.0
Mean ± SD.		28.53± 4.90
Median		28.0
Residency		
Rural	28	28.0
Urban	72	72.0
Gravidity		
Nullipara	19	19.0
Multigravida	81	81.0
Trimester		
First trimester	45	45.0
Second trimester	34	34.0
Third trimester	21	21.0

Chart 1 : Percentage of thyroid dysfunction

THYROID FUNCTION IN PREGNANCY

Table 2: Relationship between every trimester and thyroid function

Thyroid function	Trimester						x ²	MC _p
	1 st (n= 45)		2 nd (n= 34)		3 rd (n= 21)			
	No.	%	No.	%	No.	%		
Normal	23	51.1	17	50.0	11	52.4	0.848	1.000
Sub clinical hypothyroidism	17	37.8	14	41.2	8	38.1		
Clinical hypothyroidism	3	6.7	2	5.9	1	4.8		
Isolated hypothyroxinemia	2	4.4	1	2.9	1	4.8		

Table 3: Relation between thyroid function and age (years) (n= 100)

	Thyroid function								Test of sig.	P
	Normal (n= 51)		Sub clinical hypothyroidism (n= 39)		Clinical hypothyroidism (n= 6)		Isolated hypothyroidism (n= 4)			
Age (years)										
<30	30	58.8	26	66.7	3	50.0	1	25.0	x ² = 3.308	MC _p = 0.379
30-40	21	41.2	13	33.3	3	50.0	3	75.0		
Min. – Max.	18.0 – 38.0		21.0 – 39.0		22.0 – 38.0		28.0 – 35.0			
Mean ± SD.	28.55 ± 5.36		28.03 ± 4.31		29.67 ± 5.39		31.50 ± 3.11		F= 0.731	0.536
Median	28.0		28.0		29.50		31.50			

Table 4: Prevalence of thyroid dysfunction according to ATA guide-lines 2017 (n = 100)

	No.	%
Normal	77	77.0
Subclinical hypothyroidism	13	13.0
Clinical hypothyroidism	3	3.0
Isolated hypothyroxinemia	7	7.0

DISCUSSION

Considerable changes occur in the thyroid hormone physiology and thyroid gland anatomy during pregnancy. Specifically, the thyroid gland of the pregnant woman is characterized by glandular hyperplasia, increased vascularity and an approximately 30% increased volume, despite a normal echo structure^[5].

Furthermore, thyroid function test changes can occur because of estrogen-mediated increases in thyroid-binding globulin and a decline in iodide because of increased renal clearance and placental loss^[6].

In the present study, we assessed various thyroid disorders among one hundred (n=100) pregnant women attending Obstetric Outpatient Clinic in Suez-Canal University Hospital. These pregnant women ages' ranged from 18 to 39 years with average of 28.5 ± 5 years. It was found that sixty percent of the studied population was younger than thirty years old. Seventy two women were residents in urban areas. Most women were multi-gravida, representing eighty one percent. Majority of women were in their first trimester (45%) followed by second trimester (34%), while least cases were in third trimester.

This study found that most of the pregnant women (n=51) had normal thyroid functions, while subclinical hypothyroidism (n=39) was the most prevalent disorder followed by the clinical hypothyroidism (n=6) and the isolated hypothyroxinemia (n=4).

In Egypt, a cross-sectional study among 168 pregnant women who attend the Outpatient Obstetric Clinic at Ain-Shams University Hospital was carried out. The patients were subdivided into high and low risk groups for thyroid disease according to ATA guidelines as well as groups by trimester and showed that prevalence of hypothyroidism was 56% in the high risk group and 44.6% in the-low risk group with no statistically significant differences^[7].

According to these findings, the overall incidence of thyroid disorders in low risk pregnant women was 49%. This study came in accordance with who noted hypothyroidism in 55.8 % of pregnant women in a study held in Egypt mostly subclinical hypothyroidism (42.3%).

A study conducted in Jordan on 322 pregnant women in the first trimester where 20.8% of women were considered to have subclinical hypothyroidism comparable to 37.8% in this study^[8].

A cross-sectional study in Referral Hospital in Spain among 2509 pregnant women in the first trimester reported prevalence of thyroid dysfunction at 16 %^[9].

In China, a multicenter cohort study among 2899 pregnant women enrolled during their first trimester of gestation demonstrated a prevalence of thyroid dysfunction at 10.2%. This study did not use the trimester specific reference ranges recommended by American Thyroid Association^[10].

A prospective observational study in India among 1000 pregnant women attending a tertiary public hospital in the first trimester reported a prevalence of thyroid dysfunction at 14.3%, with subclinical hypothyroidism being the most common^[11].

In Sudan, cross-sectional hospital-based study among 500 pregnant Sudanese women aged 15-45 years in all trimesters, found a prevalence of 9.4 % this study use national reference ranges instead of the ATA trimester specific reference ranges^[12].

A cross-section study carried out in Tunisia among 1519 pregnant women in all trimesters demonstrated prevalence of thyroid dysfunction at 9.7%^[13].

In Saudi Arabia, across-sectional study among 384 pregnant women the majority (89.3%) of women aged less than 40 years and obesity was found to be 53.9%, the mean age was 32.6, the prevalence of subclinical hypothyroidism was 13%, and Chi-square analysis indicated that age and obesity were not significantly associated with subclinical hypothyroidism which is consistent with our results^[14].

This variability in thyroid disorders prevalence in pregnant women could be explained by the iodine status of study population, the sample size, the references used to define thyroid status or by ethnicity specificity.

Another study in Africa comparing the prevalence of thyroid functions in pregnancy using 2011 ATA guidelines and 2017 ATA guidelines. In a study by Awede *et al.*, two hundred and forty (240) pregnant women were included in the study. It showed that thyroid dysfunction was present in 24.17% of the pregnant women. Hypothyroidism was observed in 22.5% of subjects with 19.17% of subclinical hypothyroidism, 0.87% of overt hypothyroidism and 2.5% of hypothyroxinemia while hyperthyroidism was present in 1.67% of the subjects, while using 2017 ATA guidelines (TSH upper reference limit of 4.0 mU/L). In this case, thyroid disorders were

present in 14.58% of pregnant women. The frequency of subclinical hypothyroidism, overt hypothyroidism, hypothyroxinemia and hyperthyroidism were respectively 9.58%, 0.42%, 2.91% and 1.67%^[15].

In this study, similar comparison was done and revealed that the prevalence of thyroid dysfunction according to 2011 ATA reference ranges was 49% (39% SCH, 6% OH, 4% isolated hypothyroxinemia), while according to 2017 ATA reference ranges was 23% (13% SCH, 3% OH and 7% isolated hypothyroxinemia).

It was found that there was no statistically significant relationship between trimester of pregnancy and development of thyroid disorders. These results supported a study by Brent who recommended performing universal screening for thyroid dysfunction early in pregnancy as well as during the second and third trimesters^[16].

Moleti found that 40% of screened women were diagnosed with hypothyroidism in the early and late second trimester and would not have been identified, if the study had limited its observation to the first thyroid function test alone^[17].

Several researchers have also expressed great concern regarding hypothyroidism during the first trimester, as the fetus at that period is entirely dependent on maternal thyroid hormone levels, having not yet become able to make its own endogenous supply^[18].

Among different clinical parameters presented at clinics, it was found that these parameters didn't show any statistically significant relationship with certain thyroid disorders. This observation could be explained by similarity of pregnancy associated physiological changes and signs of hypo and hyperthyroidism.

Most of women were overweight (n=62), this could be related to the common belief among Egyptian pregnant women that increased caloric intake during pregnancy is encouraged for healthier fetal development. Although the weight gain in pregnancy did not correlate with hypothyroidism, it sheds lights on the importance of increasing awareness about adequate weight gain during pregnancy.

However, women with normal body mass index came next (n=25) and least cases were obese (n=13). Also, there was no statistically significant relationship between thyroid disorders and body mass index of pregnant women. These results are in contrast with Dima *et al* with a study that was done on 920 pregnant females in Lebanon found that women with a BMI > 35 kg/m² might have a greater risk of

developing hypothyroidism, this may be due to that maximum BMI in this study was 33kg/m² and also difference in sample size. They found that prevalence of hypothyroid disorders with pregnancy was 17.1% which is lower than our finding (49%)^[19].

In the current study, it was found that there was no statistically significant relationship between development of thyroid disorders and age of women, nor gravidity status nor the trimester. This comes in contrary with Awede *et al* who observed positive relationship between hypothyroidism with age, trimester and number of gravidity. This may be due to the difference in the sample size, population characteristics^[15].

CONCLUSION

In conclusion, most of the studies that were concerned about assessment of thyroid function in pregnant females noted that there was no statistically significant relationship between development of thyroid disorders and age of women, nor gravidity status.

The current study was in agreement with most of these studies. Only few studies were disagreed with these findings. Also, it was concluded that these findings from the present study underscore the prevalence of hypothyroidism among pregnant women.

CONFLICT OF INTEREST

There are no conflicts of interests.

REFERENCES

1. Jansen J, Friesema C, Milici C, Vissert J. thyroid hormone transporters in health and disease. *Thyroid* 2005 15(8):757–768.
2. Lazarus I, Brown RS, Daumerie C, Hubalewska-Dydejczyk A, Negro R, Vaidya B. European thyroid association guidelines for the management of subclinical hypothyroidism in pregnancy and in children. *Eur Thyroid J*.2014 3(2) 76-94.
3. Lazarus, J.H. *Thyroid Function in Pregnancy*. 2010 97 137-148.
4. Stagnaro-Green A, Abalovich M, Alexander E, Azizi F, Meštan J, Negro R, Nixon A, Pearce EN, Soldin OP, Sullivan S, Wiersinga W, American Thyroid Association Taskforce on Thyroid Disease During Pregnancy and Postpartum. Guidelines of the American Thyroid

- Association for the diagnosis and management of thyroid disease during pregnancy and postpartum. *Thyroid*. 2011;21(10):1081-125.
5. American College of Obstetricians and Gynecologists. Practice bulletin no. 148: thyroid disease in pregnancy. *Obstet Gynecol*. 2015;125(4):996-1005.
 6. Casey Bm, Leveno KJ. thyroid disease in pregnancy. *Obstetric Gynecology*. 2006;108:1283-1292.
 7. Ahmed IZ, Eid YM, El-Orabi H and Ibrahim HR, Comparison of universal and targeted screening for thyroid dysfunction in pregnant Egyptian women. *Eur J Endocrinol* 1, 2014 171285-291.
 8. Alkafajei A, Amarin Z, Alazaizeh W, Khader Y, Marji M. Prevalence and risk factors for hypothyroidism in Jordanian women: comparison between different reference ranges. *East Mediterr Health J*. 2012; 18(2):132–136.
 9. Dieguez M, Herrero A, Avello N, al Prevalence of Thyroid Dysfunction in Women in Early Pregnancy. 2016, 84(1):121126-.
 10. Wang W, Teng W, shanZ. The prevalence of thyroid disorders during early pregnancy in china: The benefits of universal screening in the first trimester of pregnancy. *Eur J Endocrinol* . 2011,64:263.
 11. Dhanwal DK, Prasad S, Agarwal AK . High Prevalence of subclinical hypothyroidism during first trimester of pregnancy in North India. *J EndocrMetab* . 2013,17:281-4.
 12. Saeed AK, Yassin K, Elmahdi EMA, Maralyn D, Ali AA. The Prevalence of Thyroid Dysfunction among Sudanese Pregnant Women. *SOJ GynecolObstetWomens Health* 2015; 1(1): 5.
 13. Feki M, Omar S, Menif O. Thyroid disorders in pregnancy. Frequency and association with selected diseases and obstetrical complications in Tunisian women. *ClinBiochem*. 2008,41:927-31.
 14. Al-Shanqeeti SA, Al-khudairy YN, Alabdulwahed AA, Ahmed AE, Al-Adham MS, MD, Mahmood NM, Prevalence of subclinical hypothyroidism in pregnancy in Saudi Arabia. *Saudi Med J*. 2018, 39:254-256.
 15. Awede, B., Hounnou, M., Tshabu-Aguemon, C., Adehan, G., Djrolo, F., Amoussou-Guenou, M. and Laleye, A. Thyroid Function in Pregnant Women from a West-African Population. *Open Journal of Molecular and Integrative Physiology*, 2018, 8, 1-11.
 16. Brent GA. Diagnosing thyroid dysfunction in pregnant women: is case finding enough? *Journal of Clinical Endocrinology and Metabolism*. 2007, 92 39-41.
 17. Moleti M, Lo Prešti VP, Mattina F, Mancuso A, De Vivo A, Giorgianni G, Di Bella B, Trimarchi F & Vermiglio F. Gestational thyroid function abnormalities in conditions of mild iodine deficiency: early screening versus continuous monitoring of maternal thyroid status. *European Journal of Endocrinology*. 2009, 160 611–617.
 18. Springer D, Zima T & Limanova Z. Reference intervals in evaluation of maternal thyroid function during the first trimester of pregnancy. *European Journal of Endocrinology* .2009, 160 791–797.
 19. Ezzeddine D, Hamadi C, Abbas HA, Nassar A, Abiad M, Ghazeeri G; Prevalence and Correlation of Hypothyroidism With Pregnancy Outcomes Among Lebanese Women, *Journal of the Endocrine Society*, Volume 1, Issue 5, 1 May 2017, Pages 415-422.